


**BEST AVAILABLE COPY**Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2**AMENDMENTS TO THE CLAIMS**

The following listing of claims will replace all prior versions and listing of claims in the application. For the Examiner's convenience a complete listing of all claims incorporating the amendments made herein is attached as Appendix A.

**LISTING OF CLAIMS:**

1. (Currently Amended) A process for the preparation of a compound of Formula I:



Formula I

wherein:

R<sup>1</sup> and R<sup>2</sup> are independently optionally substituted alkyl;


X is optionally substituted arylene or optionally substituted heteroarylene;

Y is a covalent bond or lower alkylene; and

Z is optionally substituted monocyclic aryl or optionally substituted monocyclic heteroaryl;

comprising;

cyclizing a compound of the formula (3):



(3)

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

wherein  $R^1$ ,  $R^2$ , X, Y, and Z are as defined above.


2. (Original) The process of claim 1, wherein the compound of formula (3) is cyclized in an inert solvent in the presence of a base.

3. (Original) The process of claim 2, wherein the inert solvent is methanol and the base is aqueous sodium hydroxide solution.

4. (Original) The process of claim 3, wherein  $R^1$  and  $R^2$  are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.


5. (Original) The process of claim 4, wherein  $R^1$  is n-propyl,  $R^2$  is ethyl, and Z is 3-trifluoromethylphenyl.

6. (Currently Amended) The process of claim 1, wherein the compound of formula (3):



(3)

is prepared by a method comprising contacting a compound of the formula (2);



(2)

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

with a compound of the formula  $Z-Y-X-CO_2H$  in the presence of a carbodiimide or with a compound of the formula  $Z-Y-X-C(O)Hal$ , where Hal is chloro or bromo.

7. (Currently Amended) The process of claim 6, wherein the compound of formula (3) is reacted with a compound of the formula  $Z-Y-X-CO_2H$  in methanol ~~in the presence of a coupling agent used to form amide bonds.~~

8. (Currently Amended) The process of claim 7, wherein the carbodiimide coupling agent used to form amide bonds is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.

9. (Currently Amended) The process of claim 6, wherein the compound of formula (3) is reacted with a compound of the formula  $Z-Y-X-C(O)Cl$ .

10. (Original) The process of claim 9, wherein the reaction is carried out in an inert solvent in the presence of a tertiary base.


11. (Original) The process of claim 10, wherein the inert solvent is acetonitrile and the tertiary base is triethylamine.

12. (Original) The process of claim 6, wherein  $R^1$  and  $R^2$  are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.

13. (Original) The process of claim 12, wherein  $R^1$  is n-propyl,  $R^2$  is ethyl, and Z is 3-trifluoromethylphenyl, namely 3-ethyl-1-propyl-8-{1-[(3-trifluoromethylphenyl)methyl]pyrazol-4-yl}-1,3,7-trihydropurine-2,6-dione.


Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2

14. (Original) The process of claim 1, wherein the compound of the formula:



(3)

is prepared by a method comprising contacting a compound of the formula;



(16)

with a compound of the formula  $R^1L$ , in which L is a leaving group.

15. (Original) The process of claim 14, wherein  $R^1$  is lower alkyl optionally substituted by cycloalkyl, and L is iodo.

16. (Original) The process of claim 15, wherein the reaction is carried out in the presence of a base in an inert solvent.


17. (Original) The process of claim 16, wherein the base is potassium carbonate and the inert solvent is N,N-dimethylformamide.

18. (Original) The process of claim 17, wherein  $R^1$  and  $R^2$  are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2


19. (Original) The process of claim 18, wherein  $R^1$  is n-propyl,  $R^2$  is ethyl, and Z is 3-trifluoromethylphenyl.

20. (Original) The process of claim 1, wherein the compound of the formula:



(3)

is prepared by a method comprising contacting a compound of the formula;



(13)

with a compound of the formula  $R^2L$ , in which L is a leaving group.

21. (Original) The process of claim 20, wherein  $R^2$  is lower alkyl optionally substituted by cycloalkyl, and L is iodo.

22. (Original) The process of claim 21, wherein the reaction is carried out in the presence of a base in an inert solvent.


23. (Original) The process of claim 22, wherein the base is potassium carbonate and the inert solvent is N,N-dimethylformamide.

Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2

24. (Original) The process of claim 23, wherein  $R^1$  and  $R^2$  are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.


25. (Original) The process of claim 24, wherein  $R^1$  is n-propyl,  $R^2$  is ethyl, and Z is 3-trifluoromethylphenyl.

26. (Currently Amended) The process of claim 14, wherein the compound of the formula:



(16)

is prepared by a method comprising contacting a compound of the formula:



(15)

with a compound of the formula  $Z-Y-X-CO_2H$  in the presence of a carbodiimide or with a compound of the formula  $Z-Y-X-C(O)Hal$ , where Hal is chloro or bromo.

27. (Currently Amended) The process of claim 26, wherein the compound of formula (15) is reacted with a compound of the formula  $Z-Y-X-CO_2H$  in methanol ~~in the presence of a coupling agent used to form amide bonds.~~

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

28. (Currently Amended) The process of claim 27, wherein the coupling-agent used to form amide bonds carbodiimide is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.

29. (Currently Amended) The process of claim 26, wherein the compound of formula (15) is reacted with a compound of the formula Z-Y-X-C(O)Cl.


30. (Original) The process of claim 29, wherein the reaction is carried out in an inert solvent in the presence of a tertiary base.

31. (Original) The process of claim 30, wherein the inert solvent is acetonitrile and the tertiary base is triethylamine.

32. (Original) The process of claim 31, wherein R<sup>1</sup> and R<sup>2</sup> are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.

33. (Original) The process of claim 32, wherein R<sup>1</sup> is n-propyl, R<sup>2</sup> is ethyl, and Z is 3-trifluoromethylphenyl.


34. (Currently Amended) The process of claim 19, wherein the compound of the formula:



(13)

is prepared by a method comprising contacting a compound of the formula:

Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2



(12)

with a compound of the formula  $Z-Y-X-CO_2H$  in the presence of a carbodiimide or a compound of the formula  $Z-Y-X-C(O)Hal$ , where Hal is chloro or bromo.

35. (Currently Amended) The process of claim 34, wherein the compound of formula (12) is reacted with a compound of the formula  $Z-Y-X-CO_2H$  in methanol in the presence of a coupling agent used to form amide bonds.

36. (Currently Amended) The process of claim 35, wherein the coupling agent used to form amide bonds carbodiimide is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.

37. (Currently Amended) The process of claim 34, wherein the compound of formula (12) is reacted with a compound of the formula  $Z-Y-X-C(O)Cl$ .

38. (Original) The process of claim 37, wherein the reaction is carried out in an inert solvent in the presence of a tertiary base.

39. (Original) The process of claim 38, wherein the inert solvent is acetonitrile and the tertiary base is triethylamine.


40. (Original) The process of claim 39, wherein  $R^1$  and  $R^2$  are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.



Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

41. (Original) The process of claim 40, wherein  $R^1$  is n-propyl,  $R^2$  is ethyl, and Z is 3-trifluoromethylphenyl.


42. (Original) The process of claim 34, wherein the compound of the formula:



(12)

is prepared by a method comprising the steps of:


a) contacting a compound of the formula:



with hexamethyldisilazane in the presence of an acid catalyst;

b) contacting the product thus formed with  $R^1L$ , where L is a leaving group, followed by;

c) contacting the product thus formed:




(10)

with a mixture of sodium nitrite in acetic acid/water; and

d) contacting the product thus formed:

Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2




(11)

with a mixture of aqueous ammonia and sodium dithionite.

43. (Original) The process of claim 42, wherein in step a) R<sup>1</sup> is lower alkyl, L is iodo, and the acid catalyst is ammonium sulfate.


44. (Original) The process of claim 26, wherein the compound of the formula:



(15)


is prepared by a method comprising the steps of:

a) contacting a compound of the formula:



with ethyl cyanoacetate in the presence of a base in a protic solvent;


b) contacting the product thus formed:



with a mixture of sodium nitrite in acetic acid/water, and

Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2


- c) contacting the product thus formed:



with a mixture of aqueous ammonia and sodium dithionite.


45. (Original) The process of claim 44, wherein the base is sodium ethoxide and the protic solvent is ethanol.

46. (Original) The process of claim 6, wherein the compound of formula:




is prepared by a method comprising the steps of:

- a) contacting a compound of the formula:



with ethyl cyanoacetate in the presence of a base in a protic solvent;


- b) contacting the product thus formed:



Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2


with the dimethylacetal of N,N-dimethylformamide;

c) contacting the product thus formed:




with a compound of formula R¹L, in which L is a leaving group;

d) contacting the product thus formed:




with aqueous ammonia;

e) contacting the product thus formed:



with a mixture of sodium nitrite in acetic acid/water; and

f) contacting the product thus formed:



with a mixture of aqueous ammonia and sodium dithionite.

47. (Original) The process of claim 46, wherein the base is sodium ethoxide and the protic solvent is ethanol.

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2


Claims 48-55 Cancelled

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

## APPENDIX B

### CLEAN COPY OF CLAIMS AS AMENDED HEREIN

1. A process for the preparation of a compound of Formula I:



Formula I

wherein:

R<sup>1</sup> and R<sup>2</sup> are independently optionally substituted alkyl;


X is optionally substituted heteroarylene;

Y is a covalent bond or lower alkylene; and

Z is optionally substituted monocyclic aryl or optionally substituted monocyclic heteroaryl;

comprising;

cyclizing a compound of the formula (3):



(3)

wherein R<sup>1</sup>, R<sup>2</sup>, X, Y, and Z are as defined above.

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2


2. The process of claim 1, wherein the compound of formula (3) is cyclized in an inert solvent in the presence of a base.

3. The process of claim 2, wherein the inert solvent is methanol and the base is aqueous sodium hydroxide solution.

4. The process of claim 3, wherein  $R^1$  and  $R^2$  are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.


5. The process of claim 4, wherein  $R^1$  is n-propyl,  $R^2$  is ethyl, and Z is 3-trifluoromethylphenyl.

6. The process of claim 1, wherein the compound of formula (3):



(3)

is prepared by a method comprising contacting a compound of the formula (2);



(2)

with a compound of the formula  $Z-Y-X-CO_2H$  in the presence of a carbodiimide or with a compound of the formula  $Z-Y-X-C(O)Hal$ , where Hal is chloro or bromo.

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

7. The process of claim 6, wherein the compound of formula (3) is reacted with a compound of the formula Z-Y-X-CO<sub>2</sub>H in methanol.

8. The process of claim 7, wherein the carbodiimide is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.

9. The process of claim 6, wherein the compound of formula (3) is reacted with a compound of the formula Z-Y-X-C(O)Cl.


10. The process of claim 9, wherein the reaction is carried out in an inert solvent in the presence of a tertiary base.

11. The process of claim 10, wherein the inert solvent is acetonitrile and the tertiary base is triethylamine.

12. The process of claim 6, wherein R<sup>1</sup> and R<sup>2</sup> are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.

13. The process of claim 12, wherein R<sup>1</sup> is n-propyl, R<sup>2</sup> is ethyl, and Z is 3-trifluoromethylphenyl, namely 3-ethyl-1-propyl-8-{1-[(3-trifluoromethylphenyl)methyl]pyrazol-4-yl}-1,3,7-trihydropurine-2,6-dione.

14. The process of claim 1, wherein the compound of the formula:




(3)

is prepared by a method comprising contacting a compound of the formula;



Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2



(16)

with a compound of the formula  $R^1L$ , in which L is a leaving group.

15. The process of claim 14, wherein  $R^1$  is lower alkyl optionally substituted by cycloalkyl, and L is iodo.


16. The process of claim 15, wherein the reaction is carried out in the presence of a base in an inert solvent.

17. The process of claim 16, wherein the base is potassium carbonate and the inert solvent is N,N-dimethylformamide.

18. The process of claim 17, wherein  $R^1$  and  $R^2$  are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.

19. The process of claim 18, wherein  $R^1$  is n-propyl,  $R^2$  is ethyl, and Z is 3-trifluoromethylphenyl.


20. The process of claim 1, wherein the compound of the formula:



Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

(3)

is prepared by a method comprising contacting a compound of the formula;




(13)

with a compound of the formula R<sup>2</sup>L, in which L is a leaving group.

21. The process of claim 20, wherein R<sup>2</sup> is lower alkyl optionally substituted by cycloalkyl, and L is iodo.
22. The process of claim 21, wherein the reaction is carried out in the presence of a base in an inert solvent.
23. The process of claim 22, wherein the base is potassium carbonate and the inert solvent is N,N-dimethylformamide.
24. The process of claim 23, wherein R<sup>1</sup> and R<sup>2</sup> are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.
25. The process of claim 24, wherein R<sup>1</sup> is n-propyl, R<sup>2</sup> is ethyl, and Z is 3-trifluoromethylphenyl.


Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

26. The process of claim 14, wherein the compound of the formula:



(16)

is prepared by a method comprising contacting a compound of the formula:



(15)

with a compound of the formula Z-Y-X-CO<sub>2</sub>H in the presence of a carbodiimide or with a compound of the formula Z-Y-X-C(O)Hal, where Hal is chloro or bromo.

27. The process of claim 26, wherein the compound of formula (15) is reacted with a compound of the formula Z-Y-X-CO<sub>2</sub>H in methanol.

28. The process of claim 27, wherein the carbodiimide is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.

29. The process of claim 26, wherein the compound of formula (15) is reacted with a compound of the formula Z-Y-X-C(O)Cl.

30. The process of claim 29, wherein the reaction is carried out in an inert solvent in the presence of a tertiary base.


Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

31. The process of claim 30, wherein the inert solvent is acetonitrile and the tertiary base is triethylamine.

32. The process of claim 31, wherein  $R^1$  and  $R^2$  are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.


33. The process of claim 32, wherein  $R^1$  is n-propyl,  $R^2$  is ethyl, and Z is 3-trifluoromethylphenyl.

34. The process of claim 19, wherein the compound of the formula:



(13)

is prepared by a method comprising contacting a compound of the formula:




(12)

with a compound of the formula  $Z-Y-X-CO_2H$  in the presence of a carbodiimide or a compound of the formula  $Z-Y-X-C(O)Hal$ , where Hal is chloro or bromo.

35. The process of claim 34, wherein the compound of formula (12) is reacted with a compound of the formula  $Z-Y-X-CO_2H$  in methanol.

Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2

36. The process of claim 35, wherein the carbodiimide is 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide.
37. The process of claim 34, wherein the compound of formula (12) is reacted with a compound of the formula Z-Y-X-C(O)Cl.
38. The process of claim 37, wherein the reaction is carried out in an inert solvent in the presence of a tertiary base.
39. The process of claim 38, wherein the inert solvent is acetonitrile and the tertiary base is triethylamine.
40. The process of claim 39, wherein R<sup>1</sup> and R<sup>2</sup> are independently lower alkyl, X is pyrazol-4-yl, Y is methylene, and Z is optionally substituted phenyl.
41. The process of claim 40, wherein R<sup>1</sup> is n-propyl, R<sup>2</sup> is ethyl, and Z is 3-trifluoromethylphenyl.
42. The process of claim 34, wherein the compound of the formula:




(12)

is prepared by a method comprising the steps of:

- a) contacting a compound of the formula:


Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2



with hexamethyldisilazane in the presence of an acid catalyst;

b) contacting the product thus formed with  $R^1L$ , where L is a leaving group, followed by;


c) contacting the product thus formed:



(10)

with a mixture of sodium nitrite in acetic acid/water; and

d) contacting the product thus formed:




(11)

with a mixture of aqueous ammonia and sodium dithionite.

43. The process of claim 42, wherein in step a)  $R^1$  is lower alkyl, L is iodo, and the acid catalyst is ammonium sulfate.


Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2

44. The process of claim 26, wherein the compound of the formula:




is prepared by a method comprising the steps of:

- a) contacting a compound of the formula:




with ethyl cyanoacetate in the presence of a base in a protic solvent;

- b) contacting the product thus formed:



with a mixture of sodium nitrite in acetic acid/water; and

- c) contacting the product thus formed:




with a mixture of aqueous ammonia and sodium dithionite.

45. The process of claim 44, wherein the base is sodium ethoxide and the protic solvent is ethanol.


Response to December 5, 2005 Office Action  
 Serial No. 10/719,102  
 CVT No. 01-0163-CIP2

46. The process of claim 6, wherein the compound of formula:




is prepared by a method comprising the steps of:

- a) contacting a compound of the formula:




with ethyl cyanoacetate in the presence of a base in a protic solvent;

- b) contacting the product thus formed:



with the dimethylacetal of N,N-dimethylformamide;

- c) contacting the product thus formed:




with a compound of formula R<sup>1</sup>L, in which L is a leaving group;

- d) contacting the product thus formed:




Response to December 5, 2005 Office Action  
Serial No. 10/719,102  
CVT No. 01-0163-CIP2




with aqueous ammonia;

e) contacting the product thus formed:



with a mixture of sodium nitrite in acetic acid/water; and

f) contacting the product thus formed:



with a mixture of aqueous ammonia and sodium dithionite.

47. The process of claim 46, wherein the base is sodium ethoxide and the protic solvent is ethanol.

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**